



Exposure of Impacted Canine

The canine, or eye tooth, normally erupts into the mouth between the ages of 11 and 13. Sometimes one or both canines develop in the wrong position. Often they lie across the roof of the mouth behind the front teeth.

Why do I need treatment?

Because one or other of your canines are in the wrong place as part of your on-going orthodontic treatment it is necessary to help the tooth erupt into the mouth. If left alone the tooth will not erupt normally and may either damage the roots of the front teeth or push them out of position.

What does the treatment involve?

Helping the tooth erupt into your mouth involves a relatively minor surgical procedure. This usually takes place under a "day case" general anaesthetic, ie although you are put to sleep completely you will be able to go home on the same day as surgery. While you are asleep the gum lying over the canine will be pushed back or a bracket (gold chain) will be glued upon the crown. Occasionally some of the bone surrounding the crown of the tooth also needs to be removed.

How will the orthodontist pull the tooth into the correct position?

Once the canine is exposed one of three things will happen under the same anaesthetic. What is going to happen for you will already have been discussed.

- **Bracket and chain.** A small bracket is glued to the tooth. Attached to this is a chain which your orthodontist can then use to pull the tooth into the right position. The chain is usually stitched out of the way but it is quite delicate and therefore it is important to be careful when eating for the first few weeks after surgery.
- **A pack.** Sometimes a pack made from gauze soaked in an antiseptic is placed over the tooth after it is exposed. The pack is kept in position with stitches and removed after a few weeks. You must be careful not to dislodge the pack. If this happens you should contact the department for advice.

Sometimes it is necessary to hold the gum back in the right position with stitches at the end of the operation. These are usually dissolvable and take about two weeks to disappear.

Is there much pain or swelling?

All of the above procedures are not particularly painful but you will obviously experience some soreness afterwards. There is usually very little in the way of swelling. If it is likely to be sore your



surgeon will arrange painkillers for you. It is not usually necessary to take antibiotics.

Will I need another appointment?

You will need to return a few weeks after surgery to have the area checked by your orthodontist or surgeon. Further appointments after this will obviously be necessary for your on-going orthodontic care.